


Definition, acronyms, nomenclature, and classification of angioedema (DANCE): AAAAI, ACAAI, ACARE, and APAAACI DANCE consensus



Avner Reshef, MD, Thomas Buttgereit, MD, Stephen D. Betschel, MD, Teresa Caballero, MD, PhD, Henriette Farkas, MD, PhD, DSc, Anete S. Grumach, MD, PhD, et al

GRAPHICAL ABSTRACT



Capsule summary: Angioedema syndromes are clinically heterogeneous, hereditary, or acquired and are caused by different pathogenetic mechanisms. Results of a global expert consensus on the classification and terminology of various angioedema types and endotypes are presented.

Definition, acronyms, nomenclature, and classification of angioedema (DANCE): AAAAI, ACAAI, ACARE, and APAAACI DANCE consensus



Avner Reshef, MD,^a Thomas Buttgereit, MD,^{b,c} Stephen D. Betschel, MD,^d Teresa Caballero, MD, PhD,^{e,mm} Henriette Farkas, MD, PhD, DSc,^f Anete S. Grumach, MD, PhD,^g Michihiro Hide, MD, PhD,^h Ankur K. Jindal, MD,ⁱ Hilary Longhurst, MD,^j Jonathan Peter, PhD,^k Marc A. Riedl, MD,^l Yuxiang Zhi, MD,^m Werner Aberer, MD,ⁿ Mohamed Abuzakouk, PhD,^o Tariq Al Farsi, MD,^p Nashat Al Sukaiti, MD,^p Mona Al-Ahmad, MD,^q Sabine Altrichter, MD,^r Emel Aygören-Pürsün, MD,^s Maria Luisa Baeza, MD, PhD,^t Noemi Anna Bara, MD,^u Andrea Bauer, MD,^v Jonathan A. Bernstein, MD,^w Isabelle Boccon-Gibod, MD,^x Hanna Bonnekoh, MD,^{b,c} Laurence Bouillet, MD, PhD,^{x,y} Zenon Brzoza, MD, PhD,^z Anette Bygum, MD, DMSc,^{aa,bb} Oscar Calderon, MD,^{cc} Regis de Albuquerque Campos, MD, PhD,^{dd} Freya Helena Campos Romero, MD,^{ee} Mauro Cancian, MD, PhD,^{ff} Herberto Jose Chong-Neto, MD, PhD,^{gg} George Christoff, MD, PhD,^{hh,ii} Stefan Cimbollek, MD,^{jj} Danny M. Cohn, MD, PhD,^{kk} Timothy Craig, DO,^{ll} Inna Danilycheva, MD,^{mmm} Razvigor Darlenski, MD, PhD,ⁿⁿ Aurélie Du-Thanh, MD, PhD,^{oo} Luis Felipe Ensina, MD, PhD,^{pp} Daria Fomina, MD,^{qq} Luz Fonacier, MD,^{rr} Atsushi Fukunaga, MD, PhD,^{ss} Asli Gelincik, MD,^{tt} Pedro Giavina-Bianchi, MD, PhD,^{uu} Kiran Godse, MD, PhD,^{vv} Mark Gompels, MD,^{ww} Margarida Goncalo, MD, PhD,^{xx} Maia Gotua, MD, PhD,^{yy} Guillermo Guidos-Fogelbach, MD, MSc, PhDc,^{zz} Mar Guilarte, MD, PhD,^{aaa} Alicja Kasperska-Zajac, MD, PhD,^{bbb} Constance H. Katelaris, MD, PhD,^{ccc} Tamar Kinaciyan, MD,^{ddd} Pavel Kolkhir, MD,^{b,c} Kanokvalai Kulthanan, MD,^{eee} Marcin Kurowski, MD, PhD,^{fff} Elena Latysheva, MD, PhD,^{mm,ggg} Antti Lauerma, MD, PhD,^{hhh} David Launay, MD, PhD,ⁱⁱⁱ Ramon Leonart, MD,^{jjj} William Lumry, MD,^{kkk} Alejandro Malbran, MD, PhD,^{lll} Ramzy Mohammed Ali, MD,^{mmm} Iman Nasr, MD,ⁿⁿⁿ Sandra Nieto-Martinez, MD, PhD,^{ooo} Claudio Parisi, MD,^{ppp} Ruby Pawankar, MD, PhD,^{qqq} Macarena Piñero-Saavedra, MD,^{rrr} Todor A. Popov, MD, PhD,^{sss} Grzegorz Porebski, MD,^{ttt} Alicia Prieto Garcia, MD, PhD,^{uuu} Polina Pyatilova, MD,^{b,c} Michael Rudenko, MD, PhD,^{vvv} Bulent Enis Sekerel, MD,^{www} Faradiba Sarquis Serpa, PhD,^{xxx} Farrukh Sheikh, MD,^{yyy} Frank Siebenhaar, MD,^{b,c} Angèle Soria, MD, PhD,^{zzz} Maria Staevska, MD, PhD,^{aaaa} Petra Staubach, MD,^{bbbb} Marcin Stobiecki, MD, PhD,^{ttt} Simon Francis Thomsen, MD, PhD,^{cccc} Massimo Triggiani, MD, PhD,^{dddd} Anna Valerieva, MD, PhD,^{aaaa} Solange Valle, MD, PhD,^{eeee} Nguyen Van Dinh, MD, PhD,^{fff} Carolina Elisa Vera Ayala, MD,^{b,c} Anna Zalewska-Janowska, MD, PhD,^{gggg} Andrea Zanichelli, MD, PhD,^{hhhh,iii} Markus Magerl, MD,^{b,c} and Marcus Maurer, MD^{b,c}

Abu Dhabi, United Arab Emirates;

Amsterdam, The Netherlands; Ankara and Istanbul, Turkey; Ashkelon, Israel; Auckland, New Zealand; Bangkok, Thailand; Barcelona, Gregorio Marañón, Madrid, and Seville, Spain; Beijing, China; Berlin, Dresden, Frankfurt, and Mainz, Germany; Bristol and London, United Kingdom; Budapest, Hungary; Buenos Aires, Argentina; Campbelltown, Australia; Cape Town, South Africa; Chandigarh and Maharashtra, India; Cincinnati, Ohio; Coimbra, Portugal; Copenhagen and Odense, Denmark; Curitiba, Espirito Santo, Salvador, Santo André, São Paulo, and Rio De Janeiro, Brazil; Dallas, Tex; Doha, Qatar; Graz, Linz, and Vienna, Austria; Grenoble, Lille, Montpellier, and Paris, France; Hanoi, Vietnam; Helsinki, Finland; Hershey, Pa; Hiroshima and Osaka, Japan; Katowice, Lodz, and Opole, Poland; Kuwait City, Kuwait; La Jolla, Calif; Lahore, Pakistan; Lima, Peru; Mexico City, Mexico; Milan, Padua, and Salerno, Italy; Mineola, NY; Moscow, Russia; Muscat, Oman; Sângeorgiu de Mureș, Romania; Sofia and Stara Zagora, Bulgaria; Tbilisi, Georgia; and Toronto, Ontario, Canada

From ^athe Angioedema Research Center, Barzilai University Medical Center, Ashkelon; ^bthe Angioedema Center of Reference and Excellence (ACARE), Institute of Allergology, Charité—Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin; ^cthe Fraunhofer Institute for Translational Medicine, and Pharmacology ITMP, Immunology and Allergology, Berlin; ^dthe Department of Medicine, St Michael's Hospital, University of Toronto, Toronto; ^ethe Department of Allergy, La Paz University Hospital, Hospital La Paz Institute for Health Research (IdiPAZ—Group 44), Biomedical Research Network on Rare Diseases (CIBERER U754), Madrid; ^fthe Hungarian Angioedema Center of Reference and Excellence, Department of Internal Medicine and Haematology, Semmelweis University, Budapest; ^gthe Department of Clinical Immunology, University Center Faculdade de Medicina do ABC, Santo André; ^hthe Department of Dermatology, Institute of Biomedical & Health Sciences, Hiroshima University, Hiroshima; ⁱthe Department of Pediatrics, Allergy Immunology Unit, Postgraduate Institute of

Medical Education and Research, Chandigarh; ^jthe Department of Medicine, University of Auckland and Department of Immunology, Auckland City Hospital, Auckland; ^kthe Department of Medicine, Division of Allergy and Clinical Immunology, University of Cape Town, and the Allergy and Immunology Unit, University of Cape Town Lung Institute, Cape Town; ^lthe Division of Rheumatology, Allergy, and Immunology, University of California San Diego, La Jolla; ^mthe Department of Allergy, Peking Union Medical College Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing; ⁿthe Department of Dermatology, Medical University of Graz, Graz; ^othe Allergy and Immunology, Respiratory Institute, Cleveland Clinic Abu Dhabi, Abu Dhabi; ^pthe Department of Pediatric Allergy and Clinical Immunology, The Royal Hospital, Muscat; ^qthe Microbiology Department, College of Medicine, Kuwait University, Kuwait City; ^rthe Klinik für Dermatologie und Venerologie, Kepler Uniklinikum, Linz; ^sthe Department of Pediatrics, University Hospital Frankfurt, Goethe University, Frankfurt; ^tthe Allergy Department, Hospital General

Background: Angioedema (AE) manifests with intermittent, localized, self-limiting swelling of the subcutaneous and/or submucosal tissue. AE is heterogeneous, can be hereditary or acquired, may occur only once or be recurrent, may exhibit wheals or not, and may be due to mast cell mediators, bradykinin, or other mechanisms. Several different taxonomic systems are currently used, making it difficult to compare the results of studies, develop multicenter collaboration, and harmonize AE treatment.

Objective: We developed a consensus on the definition, acronyms, nomenclature, and classification of AE (DANCE).

Methods: The initiative involved 91 experts from 35 countries and was endorsed by 53 scientific and medical societies, and patient organizations. A consensus was reached by online discussion and voting using the Delphi process over a period of 16 months (June 2021 to November 2022).

Results: The DANCE initiative resulted in an international consensus on the definition, classification, and terminology of AE. The new consensus classification features 5 types and endotypes of AE and a harmonized vocabulary of abbreviations/acronyms.

Conclusion: The DANCE classification complements current clinical guidelines and expert consensus recommendations on

the diagnostic assessment and treatment of AE. DANCE does not replace current clinical guidelines, and expert consensus algorithms and should not be misconstrued in a way that affects reimbursement of medicines prescribed by physicians using sound clinical judgment. We anticipate that this new AE taxonomy and nomenclature will harmonize and facilitate AE research and clinical studies, thereby improving patient care. (J Allergy Clin Immunol 2024;154:398-411.)

Key words: Angioedema, acronyms, classification, terminology, types

Angioedema (AE) is defined as “intermittent, localized, and self-limiting swelling of the subcutaneous and/or submucosal tissue, due to a temporary increase in vascular permeability.”^{1,2} The pathophysiologic abnormalities in AE are attributed to mechanisms controlling endothelial barrier function, which normally maintain the stability of capillaries and small venules.³ In AE, dysregulated endothelial permeability leads to excess plasma leakage from the blood vessels into the surrounding tissues, resulting in transient swelling.³ AE manifests with recurrent, reversible swelling episodes of skin and subcutaneous tissues or

Universitario Gregorio Marañón, Biomedical Research Network on Rare Diseases—U761, Gregorio Marañón Health Research Institute (IISGM), Gregorio Marañón, Madrid; ^uthe Romanian Hereditary Angioedema Expertise Centre, Centrul Clinic Mediu-quest, Sângeriu de Mureș; ^vthe Department of Dermatology, University Hospital Carl Gustav Carus, Technical University Dresden, Dresden; ^wthe University of Cincinnati College of Medicine, Department of Internal Medicine, Division of Rheumatology, Allergy and Immunology, Cincinnati; ^xthe National Reference Center of Angioedema CREAK, Grenoble; ^ythe Internal medicine department, Grenoble University Hospital, Grenoble; ^zthe Department of Internal Diseases with Division of Allergy, Institute of Medical Sciences, University of Opole, Opole; ^{aa}the Department of Clinical Genetics, Odense University Hospital, Odense; ^{bb}the Clinical Institute, University of Southern Denmark, Odense; ^{cc}Clínica SANNA el Golf, San Isidro, Lima; ^{dd}the Universidade Federal da Bahia, Brazilian Network of Hereditary Angioedema, Salvador; ^{ee}the Department of Allergy and Clinical Immunology, Hospital Central Sur Alta Especialidad, Mexico City, Mexico City; ^{ff}the Departmental Unit of Allergology, University Hospital of Padua, Padua; ^{gg}the Serviço de Alergia e Imunologia, Complexo Hospital de Clínicas, Universidade Federal do Paraná, Curitiba; ^{hh}Excelsior Medical Centre, Sofia; ⁱⁱMedical University—Sofia, Sofia; ^{jj}the Hospital Universitario Virgen del Rocío, Seville; ^{kk}the Department of Vascular Medicine, Amsterdam UMC, Amsterdam Cardiovascular Sciences, University of Amsterdam, Amsterdam; ^{ll}the Department of Pediatrics, Pennsylvania State University, Hershey; ^{mm}the NRC Institute of Immunology FMBA of Russia, Moscow; ⁿⁿthe Department of Dermatovenereology, Trakia University, Stara Zagora; ^{oo}the Département de dermatologie, ACARE, Centre Hospitalier Universitaire de Montpellier; Montpellier; ^{pp}the Federal University of São Paulo, São Paulo; ^{qq}the Moscow City Research and Practical Center of Allergology and Immunology, Clinical Hospital No. 52, Moscow Healthcare Department, Moscow; ^{rr}the New York University—Long Island School of Medicine, Mineola; ^{ss}the Department of Dermatology, Division of Medicine for Function and Morphology of Sensory Organs, Faculty of Medicine, Osaka Medical and Pharmaceutical University, Osaka; ^{tt}the Division of Immunology and Allergic Diseases, Department of Internal Medicine, Istanbul Faculty of Medicine, Istanbul University, Istanbul; ^{uu}the Division of Clinical Immunology and Allergy, Universidade de São Paulo, São Paulo, São Paulo; ^{vv}the Dr D. Y. Patil Medical College and Hospital, Navi Mumbai, Maharashtra; ^{ww}the Department of Immunology, ACARE, North Bristol NHS Trust, Bristol; ^{xx}the Department of Dermatology, Coimbra Hospital and University Center, and Faculty of Medicine, University of Coimbra, Coimbra; ^{yy}the Center of Allergy and Immunology and David Tivdiani Medical University, Tbilisi; ^{zz}the SEPI/ENMH Instituto Politécnico Nacional, Mexico City; ^{aaa}the Allergy Department, Hospital Universitario Vall d’Hebron, Vall d’Hebron Research Institute (VHIR), Barcelona; ^{bbb}the European Center for Diagnosis and Treatment of Urticaria and Angioedema and Department of Clinical Allergy and Urticaria, Medical University of Silesia in Katowice, Katowice; ^{ccc}Western Sydney University and Campbelltown Hospital, Campbelltown; ^{ddd}the Department of Dermatology, Medical University of Vienna, Vienna; ^{eee}the Department of Dermatology, Faculty of Medicine Siriraj Hospital, Mahidol University, Bangkok; ^{fff}the Department of Immunology and Allergy, Medical University of Lodz, Lodz; ^{ggg}Pirogov Russian National Research Medical University, Moscow; ^{hhh}the Department of Dermatology, Skin and Allergy Hospital, Helsinki University Hospital,

Helsinki; ⁱⁱⁱUniversity Lille, Inserm, CHU Lille, Service de Médecine Interne et Immunologie Clinique, Centre de Référence des Angioedemes à Kinine (CREAK), U1286-INFINITE—Institute for Translational Research in Inflammation, Lille; ^{jjj}the Allergology Department, Hospital Universitari Bellvitge, IDIBILL Research Institute, L’Hospitalet de Llobregat, Barcelona; ^{kkk}the AARA Research Center, Dallas; ^{lll}the Unidad de Alergia, Asma e Inmunología Clínica, Buenos Aires; ^{mmm}the Department of Medicine, Allergy and Immunology Division, Hamad Medical Corporation, Doha; ⁿⁿⁿthe Immunology and Allergy Department, The Royal Hospital, Muscat; ^{ooo}the Unidad de Genética de la Nutrición, Instituto Nacional de Pediatría, Mexico City; ^{ppp}the Pediatric and Adult Allergy sections of the Italian Hospital of Buenos Aires, Buenos Aires; ^{qqq}Nippon Medical School, Tokyo, Japan; ^{rrr}the Allergy Department, Centro Hospitalar do Algarve, Faro, Portugal; ^{sss}University Hospital Sv Ivan Rilski, Sofia; ^{ttt}the Department of Clinical and Environmental Allergology, Jagiellonian University Medical College, Krakow, Poland; ^{uuu}the Department of Allergy, Gregorio Marañón Health Research Institute (IISGM), Gregorio Marañón University Hospital, Madrid; ^{vvv}The London Allergy and Immunology Centre, London; ^{www}the Pediatric Allergy and Asthma Division, Hacettepe University, Ankara; ^{xxx}the Angioedema and Urticaria Reference Center, Hospital Santa Casa de Misericórdia de Vitória, Vitória, Espírito Santo; ^{yyy}the Chughtai Medical Center, Lahore; ^{zzz}the Médecine Sorbonne Université, Service de Dermatologie et Allergologie, hôpital Tenon, Assistance Publique Hôpitaux de Paris, Paris; ^{aaaa}the Department of Allergology, Medical University of Sofia, Clinic of Allergology, University Hospital “Alexandrovska,” Sofia; ^{bbbbb}the Department of Dermatology, ACARE, University Medical Center Mainz, Mainz; ^{ccccc}the Department of Dermatovenereology, Bispebjerg Hospital, Copenhagen; ^{dddd}the Division of Allergy and Clinical Immunology, University of Salerno, Salerno; ^{eeeee}the Department of Internal Medicine, Immunology Service, Hospital Universitario Clementino Fraga Filho, Rio De Janeiro; ^{ffff}the Department of General Internal Medicine, Respiratory—Allergy and Clinical Immunology Unit, Vinmec Healthcare System, Hanoi; ^{gggg}Department of Psychodermatology, Medical University of Lodz, Lodz; ^{hhhhh}Department of Biomedical Sciences for Health, Università degli Studi di Milano, Milan; and ⁱⁱⁱⁱthe Operative Unit of Medicine, IRCCS Policlinico San Donato, San Donato Milanese, Milan.

The first 2 authors contributed equally to this article, and both should be considered first author. The last 2 authors contributed equally to this article, and both should be considered senior author.

Received for publication November 28, 2023; revised February 24, 2024; accepted for publication March 4, 2024.

Available online April 25, 2024.

Corresponding author: Avner Reshef, MD, Angioedema Research Center, Barzilai University Medical Center, Ashkelon, Israel. E-mail: areshef@netvision.net.il.

 The CrossMark symbol notifies online readers when updates have been made to the article such as errata or minor corrections

0091-6749/\$36.00

© 2024 Published by Elsevier Inc. on behalf of the American Academy of Allergy, Asthma & Immunology

<https://doi.org/10.1016/j.jaci.2024.03.024>

submucosal surfaces. Involvement of the upper airway is life-threatening, and intestinal wall swelling may result in obstruction and ascites.^{4,5}

Most recurrent AE episodes are driven by mast cell (MC) mediators, as in urticaria (URT), a disease defined by the international guideline for URT as the occurrence of wheals, AE, or both.⁶ In chronic spontaneous URT (CSU), most patients report recurrent wheals (hives), often with concomitant AE, but in ~10%, AE occurs without wheals.^{6,7}

Hereditary AE (HAE) due to C1 inhibitor (*C1INH*) deficiency is a rare genetic disorder resulting from low levels or dysfunctional *C1INH*, the primary role of which is to control blood vessel permeability.⁸ HAE due to *C1INH* deficiency was first described by Donaldson and Evans,⁹ followed by a series of clinical cases by Landerman.¹⁰ Over the last 57 years, more than 850 different *C1INH* gene (*SERPING1*) mutations have been documented, located on chromosome 11.¹¹⁻¹³ Misdiagnosis of intestinal swelling may lead to unnecessary abdominal surgery,⁵ and attacks involving the oropharyngeal–lingual region, if not promptly recognized and treated, may cause death due to asphyxiation.^{14,15} As a lifetime chronic illness, HAE severely affects all aspects of daily activities and health-related quality of life, and it considerably burdens health care systems.¹⁶⁻¹⁹ International guidelines regarding the diagnosis and management of HAE have been published and regularly updated.²⁰⁻²²

Recent evidence shows that the mechanisms underlying AEs are heterogeneous and complex. Examples include the disinhibition of the kallikrein/kinin system (KKS), leading to local and transient excess production of bradykinin (BK), as in hereditary or acquired types of *C1INH* deficiency, reduced catabolism of BK (ie, drug induced), and intrinsic abnormalities in elements of the vascular endothelium.

In the last 2 decades, several new HAE syndromes were described, all with normal quantity and activity of *C1INH* cosegregating with different gene mutations.²³⁻²⁹ In view of the aforementioned new evidence, the conventional paradigm of AE classification needs to be reevaluated.³⁰

Previous classifications of AE

Several attempts to classify AE have been made in the past. An extensive literature search in 2016 found 25 international consensus statements on the definition and classification of AE published over the previous 13 years.³¹ The pivotal consensus statement by Cicardi et al¹ provided an extensive evidence-based foundation for all forms of AE known at that time, including incidence, clinical presentation, pathogenesis, diagnosis, and treatment. Subsequent consensus statements, guidelines, and practice parameters provided updated and revised classifications of AE.²⁰⁻²²

Rationale for updated classification and terminology of AE

Currently, AE terminology and abbreviations/acronyms in the medical literature differ by country, professional organizations, and authors. Consequently, dissimilar wording and terminology has led to confusion and inconsistencies.³¹ The Hereditary Angioedema Working International Group (HAWK) consensus was the first to formulate an AE-specific vocabulary of acronyms

and taxonomy.¹ However, the diverse vocabulary used in publications motivated a call for unified terminology.³²⁻³⁴

The current AE classification system leaves many taxonomical gaps. The finding of novel mutations causing HAE led to the general term “HAE with normal C1 inhibitor,” based on the presence or absence of *C1INH* during a preliminary assessment. Also, MC-mediated AEs (ie, CSU) are separately classified, and other common AEs are not yet listed by any classification system.⁶ In addition, AE due to drugs (ie, angiotensin-converting enzyme [ACE] inhibitors) and rare AE diseases caused by vascular hyperpermeability are also not included.

These shortcomings call for an updated, comprehensive, and widely accepted classification system, one that combines clinical expression, mechanisms, biomarkers, and genetics, and that can reflect all the different AE types. Such a classification system will harmonize publications’ vocabulary and improve patient care by avoiding misdiagnosis and assisting in tailoring targeted therapies.

The definition, acronyms, nomenclature, and classification of AE (DANCE) initiative addresses this unmet need and provides a global consensus on the definition and classification of AE and its nomenclature, including abbreviations/acronyms, covering all AE types and subtypes.

METHODS

Literature search

On June 22, 2021, two independent reviewers (T.B. and C.V.) separately conducted a systematic PubMed literature search of the last 10 years with the MeSH terms “angioedema AND definition” (76 results) and “angioedema AND classification” (178 results), in accordance with PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analysis) guidelines.³⁵ All studies were published in German, English, or Spanish. Additionally, we included and manually checked references in key publications on AE because definitions and classifications of AE are often only reported as part of the patient or study description in the full text.

The studies had to report a definition and/or classification of AE to be eligible. On the basis of these predefined criteria, the reviewers screened all study titles and abstracts for suitability and subsequently reviewed suitable articles in full text to evaluate their eligibility. Finally, 44 publications were included and analyzed.

The following details were extracted from each study: first author’s name and year of publication; definition of AE; classification of AE; and names and abbreviations of AE types and subtypes. This information was made available to the DANCE steering committee and expert panel members.

DANCE steering committee and expert panel

The DANCE steering committee consisted of 13 AE experts, diverse in both professional background (8 clinical immunologists/allergists, 3 dermatologists, 1 pediatrician) and geography (Brazil, Canada, China, Germany, Hungary, India, Israel, Japan, New Zealand, South Africa, Spain, United States). The steering committee was led by Avner Reshef (Israel), Markus Magerl (Germany), and Marcus Maurer (Germany).

The DANCE expert panel initially recruited 97 experts. Five experts did not respond in round 1, and 1 did not respond in round 3. Consequently, 91 experts from 35 countries, including the

steering committee members, contributed to the consensus. They came from different fields of expertise, including dermatology, allergy, immunology, pediatrics, otolaryngology, gastroenterology, and emergency medicine, as delegates of 53 medical societies (see Table E1 in the Online Repository available at www.jacionline.org). Participation in the expert panel was facilitated by publicized calls and a registration platform on the Angioedema Centers of Reference and Excellence (ACARE) website to represent interdisciplinary and intercontinental views on AE and to gain a robust consensus (acare-network.com/projects/#past). All expert panel members confirmed that they are actively treating patients with AE and/or were involved in research directly related to AE, and that they had obtained a mandate to be the delegate of a national or international scientific or medical society; they further confirmed in writing their nomination of experts as delegates, endorsement of the outcome of this project, and support of its dissemination. No honoraria were paid for participation.

Development of rationale and concept of DANCE

On the basis of the literature search results, the steering committee leaders developed the rationale and general concept for the DANCE initiative. The first draft set of statements was presented to the steering committee in video conferences in September and October 2021. After the group had agreed on the scope of the initiative and the rationale and principles of the new classification, the steering committee leaders continued to refine the definition of AE, names assigned to the different types and subtypes of AE, abbreviations/acronyms, and classification. After further discussion, 18 statements on the nomenclature and classification of AE were proposed and presented to the panel members for the first round of voting, using a Delphi process (Fig 1). The proposed nomenclature reflected the current International Classification of Diseases and Related Health Problems (ICD) and the therapeutic indications and target diseases used in the summaries of product specifics of current treatments for AE.

Establishment of consensus by Delphi process

The global expert panel established the core consensus via a Delphi process over a period of 8 months (March 14 to November 15, 2022). The Delphi process is a validated approach to evaluate and refine group opinions (*vox populi*) by iterative rounds of questioning without the influence of group pressure or dominant individuals.³⁶ The process enables a free exchange of views while ensuring that all opinions are equally considered.³⁶⁻³⁸ The DANCE Delphi process consisted of 3 rounds of voting, handled by Thomas Buttgeriet, a trained Delphi facilitator. A dedicated online polling system (Welphi, Decisioneyes, Lisbon, Portugal) was used, similar to a previous Delphi process on HAE treatment goals.³⁹

In round 1 (49 days, from March 14 to May 1, 2022, with 92 respondents) of the DANCE Delphi process, free-text responses of the expert panel members to the suggestions of the steering committee for the definition, nomenclature, and classification for AE were collected, together with the evaluation of each suggestion by use of a 5-point Likert scale (0 = completely disagree, 1 = mostly disagree, 2 = undecided, 3 = mostly agree, 4 = completely agree). Respondents were asked to consider their clinical experience, their patient management protocols, and their broader knowledge. Consensus was defined *a priori* as agreement

(mostly agree or completely agree) by at least 75% of respondents. Percentage agreement and all comments were recorded for each statement.³⁸ The Delphi facilitator provided an anonymized summary of the outcomes and presented the results to the panel members. The steering committee developed a revised version of the statements that was based on the results and feedback obtained in round 1.

In round 2 (from July 21 to August 14, 2022), expert panel members were asked if they agreed or disagreed with the revised statements. After round 2, consensus was reached on the definition of AE and 15 of 18 additional statements, and the outcomes were again summarized and presented to the expert panel members. For the third and final round (from October 25 to November 15, 2022), the steering committee developed revised versions of the 3 statements for which consensus was not achieved in round 2. These revised statements considered the feedback of panel members who did not agree during round 2, with additional information, evidence, and context wording. The third Delphi round established consensus on all revised statements (Fig 1 and Table I).

RESULTS

Definition of AE and related terms

The present DANCE consensus provides 3 AE-related definitions (Table I, statements 1-3). The consensus definition of AE is “a paroxysmal, localized, and self-limiting swelling of the subcutaneous and/or submucosal tissue, due to a temporary increase in vascular permeability.”¹ This definition highlights the 3 key features of manifestations of AE. The panel preferred “paroxysmal” to “intermittent” to emphasize the acute nature of AE attacks. Because swelling can simultaneously affect subcutaneous and submucosal tissues, the wording “and/or” was chosen. The DANCE expert panel chose not to include “potentially life-threatening” because death from asphyxiation is rare and does not occur in all types of AE.

The second definition (Table I, statement 2) relates to wheals (hives). These welts occur mainly in URT and are characterized by 3 typical features: a sharply circumscribed superficial central swelling of variable size and shape, almost invariably surrounded by reflex erythema; an itching or sometimes burning sensation; and its fleeting nature, with the skin returning to its normal appearance, usually within 30 minutes to 24 hours, as defined by a multidisciplinary guideline.⁶ In a recent systematic review, two-thirds of patients with CSU were reported to have concomitant AE in one study, and the pooled prevalence of AE across all CSU studies analyzed was 36.5%.⁴⁰ Some experts thought this definition should not emphasize the evanescent nature of the wheals because they can persist for several days. Patients with wheals that last longer than 24 hours should be assessed for other diseases, including urticarial vasculitis.

Our third definition (Table I, statement 3) relates to AE prodromes. These signs or symptoms (1) predate and often indicate the onset of an AE attack, (2) manifest subjectively (ie, pain, tingling, anxiety) or objectively (ie, skin rash, flushing, diarrhea), (3) typically occur 1 to 12 hours before the onset of an AE attack, and (4) may recur in the same manner, allowing patients to predict an impending attack.⁴¹ This definition is based on many observations, clinical studies, and publications on prodromes as predictors of AE attacks in patients with HAE.^{41,42} Some experts commented that there is a need for a clearer demarcation of the




FIG 1. Timeline of the Delphi process (May 2021 to November 2022). Experts from 35 countries, including steering committee members, representing 53 medical societies, contributed to the consensus. More information is provided in Tables E5 and E6 in the Online Repository available at www.jacionline.org.

time course between prodromes and attacks regarding location and clinical features. Nonetheless, recent studies show that HAE patients can distinguish prodromes from attacks, and prodromes can foretell subsequent attack location and intensity.⁴¹ Because prodromes are not necessarily followed by an attack, the words “often” and “possible” were introduced to indicate that prodromes are not always followed by an AE attack.

Nomenclature of AE

The DANCE initiative recommends using “AE” as the abbreviation for all types of AE (Table I, statement 4). This standard, etymology-driven, and short term uses the first letters of the 2 components of the word “angioedema.” However “AE” also serves to abbreviate other terms, including “adverse events” in clinical trials. When used in publications, its introduction at first use and application in reference to angioedema will minimize confusion. Some experts suggested including an abbreviation for chronic or recurrent AE. However, the minimum number of attacks, or time during which swellings occur in recurrent or chronic AE, is not well defined and requires further research.

The DANCE expert panel further recommends that abbreviations/acronyms for inherited AE types start with the letter H, for hereditary (ie, HAE), and that the letter A, for acquired, is reserved for the abbreviation of AE due to acquired C1 inhibitor deficiency, traditionally abbreviated AAE (Table I, statement 5). Most cases and types of AE are acquired rather than hereditary; therefore, “acquired” is not needed for the names of nonhereditary subtypes of AE. The exception to this rule is AE due to acquired *C1INH* deficiency (AAE-C1INH). As a result of the wide overlap between

different etiologies of acquired *C1INH* deficiency, DANCE’s classification follows the previous classification consensus and does not divide AE-C1INH into subcategories.¹

Abbreviations/acronyms of AE subtypes are recommended to start with the disease first (ie, HAE, AAE, AE), followed by a dash/hyphen and mechanism or biomarker (Table I, statement 6). This approach differs from previous ones, including the 2014 HAWK group nomenclature.¹ The panel agreed that starting the designation of AE subtypes with the clinical presentation (phenotype) provides the most crucial information first and whether it is hereditary or acquired. Indication of the underlying pathomechanism or the associated biomarker then follows. This will simplify abbreviations of AE subtypes (Table I, statement 7).

Two forms of HAE due to *C1INH* deficiency are described: HAE due to *C1INH* quantitative deficiency (characterized by low antigenic and functional *C1INH* levels) and HAE due to *C1INH* dysfunction (characterized by normal or elevated antigenic but low functional *C1INH* levels). These subtypes are traditionally referred to as type 1 and type 2, respectively. Thus, the panel recommends that they be abbreviated HAE-C1INH-Type1 and HAE-C1INH-Type2, respectively (Table I, statement 8). This intends to preserve this established HAE terminology and align it with recent practice parameters.²⁰⁻²²

In addition, the abbreviations of coagulation-contact system components proposed by Schmaier et al⁴³ were adopted (Tables I and II, statement 9). This publication, authored by 11 distinguished hematology experts, was approved by the International Society on Thrombosis and Hemostasis.⁴³ The authors recommended that the new abbreviations be utilized in all future communications and publications in the field.

TABLE I. Summary of key statements and rate of agreement obtained after 3 rounds of Delphi process

Statement no.	Statement	Agreement*
1	Angioedema is defined as “a paroxysmal, localized, and self-limiting swelling of the subcutaneous and/or submucosal tissue, due to a temporary increase in vascular permeability” (modified from Cicardi et al, ¹ Allergy 2014).	98%
2	Wheal in urticaria is characterized by 3 typical features (modified from Zuberbier et al, ⁶ Allergy 2022): 1. Sharply circumscribed superficial central swelling of variable size and shape, almost invariably surrounded by reflex erythema. 2. Itching or sometimes burning sensation. 3. Fleeting nature, with skin returning to normal appearance, usually within 30 minutes to 24 hours.	99%
3	Prodromes of AE are characterized by (modified from Leibovich-Nassi and Reshef, ⁴¹ Clin Rev Allergy Immunol 2021): 1. Sign or symptom often indicating impending onset of attack. 2. Manifested subjectively (ie, pain, tingling, anxiety) or objectively (ie, skin rash, flushing, diarrhea). 3. Typically occur 1 to 12 hours before attack. 4. May reiterate in the same manner, allowing patients to predict possible onset of upcoming attacks.	97%
4	The collective acronym for all types of angioedema will be AE (plural, AEs).	100%
5	Acronyms for inherited angioedema (AE) types will start with letter H, for hereditary (ie, HAE-C1INH); and use of letter A, for acquired, before AE is reserved for AE due to acquired C1INH deficiency (ie, AAE-C1INH).	99%
6	Acronyms will assemble by using disease first (ie, HAE), followed by mechanism/biomarker, which may be separated by dash or hyphen (ie, HAE-FXII, AE-ACEI).	99%
7	Mechanisms, mediators, or biomarkers will be inscribed without dash or hyphen (ie, ACEI, C1INH).	95%
8	The 2 known types of HAE due to C1INH deficiency will use suffix Type1 or Type2 (ie, HAE-C1INH-Type1, HAE-C1INH-Type2).	99%
9	Acronyms of all coagulation-contact system components will follow definitions of Schmaier et al ⁴³ (J Thromb Haemost 2019).	99%
10	Acronyms currently used to describe AE treatment strategies will remain the same (ie, ODT, STP, LTP).	99%
11	AE categories will be assembled into 3 endotypes and 2 subgroups.	89%
12	All recurrent mast cell-mediated AEs (histamine mediated, “allergic anaphylactic”) will be designated as AE-MC endotype.	99%
13	All AEs related to pathomechanism of the contact kallikrein/kinin system (KKS) will be designated as AE-BK endotype.	98%
14	All AEs related to intrinsic vascular endothelium abnormalities and mutations will be designated as AE-VE endotype.	97%
15	HAE-HS (HS3ST6 mutation) is categorized as AE-VE endotype.	98%
16	All drug-induced AEs without proven pathomechanisms will be designated as AE-DI type.	85%
17	ACE inhibitor-induced AE is categorized as AE-DI type.	93%
18	AEs with still-unknown mechanism/triggers will be designated as AE-UNK type.	90%
19	Episodic AE with eosinophilia (EAE, Gleich syndrome) is categorized as AE-UNK type.	100%

FXII, Factor XII; LTP, long-term prophylaxis; ODT, on-demand treatment; STP, short-term prophylaxis.

*Rate of agreement reflects percentage of experts who agreed with the statement, as determined during 3 Delphi rounds of consensus finding.

The DANCE experts propose that the abbreviations currently used to describe AE treatment strategies remain the same—that is, ODT, STP, and LTP for on-demand treatment, short-term prophylaxis, and long-term prophylaxis, respectively (Table I, statement 10). These terms and their abbreviations already have their place in the AE literature, clinical trials, and practice parameters and are well established in HAE treatment plans.

Classification of AE

AE subtypes will be assigned to 5 different types, the first 3 of which designate AE endotypes (Table I, statement 11, Fig 2). The DANCE classification is meant to complement current guidelines and other consensus algorithms for the diagnosis and treatment of AE patients. It aims to improve the characterization of AE subtypes so that ongoing efforts to investigate mechanisms of these poorly understood conditions will lead to more meaningful and generalizable results.²⁰⁻²² The DANCE classification of AE uses all the currently available data to formulate terms that reflect phenotypic expressions, genetics, mechanisms, and biomarkers but acknowledges that specific biomarkers are still lacking for

TABLE II. Abbreviations/acronyms of coagulation-contact system components

Name	Acronym
Bradykinin	BK
Cleaved high-molecular-weight kininogen	cHK
Activated factor XII	FXIIa
Coagulation factor XII (Hageman factor)	FXII
Coagulation factor XI (Rosenthal factor)	FXI
High-molecular-weight kininogen (Williams/Fitzgerald factor)	HK
Low-molecular-weight kininogen	LK
Plasma prekallikrein (Fletcher factor)	PK
Plasma kallikrein	PKa

Modified from Schmaier et al⁴³ (J Thromb Haemost 2019).

many conditions, so changes may be needed as new evidence emerges. Any of the proposed categorizations may change as more evidence becomes available. It is important to emphasize that the DANCE classification and its future updates should not be misconstrued in a way that would affect reimbursement of physician-prescribed medications for HAE variants.




FIG 2. Proposed new classification of all AE entities according to clinical expression, currently available information on pathomechanisms, and known pathogenesis. Five types/endotypes are listed by their putative mechanisms and new terminology (abbreviation/acronym). The new classification should not be used as a barrier to the medical treatment of patients with HAE with normal C1INH when based on a clinician's judgment. HAE-UNK (ie, HAE due to unknown mutation) can be bradykinin-mediated as evidenced by the response to bradykinin-targeted treatment. It can also be vascular endothelium dysfunction-mediated, and it can be truly unknown in its etiology and mechanism. See Table III for the meanings of abbreviations/acronyms used in this illustration.

MC-mediated AE

Mast cell-mediated AE (AE-MC) replaces “histamine-mediated AE,” “allergic,” or “anaphylactic”; and will be considered an endotype (Table I, statement 12). This change is based on the understanding that MCs secrete many proinflammatory products beyond histamine.⁴⁴ Thus, this endotype is based on a cell rather than a mediator, and recurrent AE-MC (ie, CSU) often does not respond to higher-than-standard doses of antihistamines, although it often does respond to omalizumab. Subtypes of AE-MC include acute URT, inducible URT, and CSU, with up to 70% of patients developing AE.^{6,7} AE-MC also comprises AE in the context of anaphylaxis because it often manifests with AE attacks at various organs and carries the risk of asphyxiation if involving the oropharyngeal area.⁴⁵

BK-mediated AE

The AE-BK endotype comprises all BK-mediated subtypes of AE (Table I, statement 12), including all AEs attributed to the KKS cascade and BK production.⁸ Of note, all types of AE attributed to the activation of the KKS are held to be BK driven, but BK may be generated by KKS-independent mechanisms in some types of AE—for example, in HAE-PLG—by a mutant plasmin (plasmin-Glu³¹¹).⁴⁶ In addition to including HAE-C1INH and AAE-C1INH, which are both associated with C1INH deficiency, this category will comprise 3 recently identified genetic variants, namely HAE-FXII, HAE-PLG, and HAE-KNG, as well as HAE-UNK (unknown).

The DANCE classification distinguishes between hereditary and acquired AE, which comprise the known conditions caused by C1INH deficiency. All types of HAE that are not caused by C1INH deficiency are clinically categorized as HAE-nC1INH (Fig 3), with further subcategorization by the genetic or molecular cause—for example, HAE-nC1INH due to factor XII (FXII) mutation.

AE due to intrinsic vascular endothelium dysfunction

This new category refers to intrinsic vascular endothelium dysfunction as an endotype (Table I, statement 14). The decision to form a new category within this updated classification is based on new evidence on the central role of the vascular endothelium in the pathologic processes leading to AE attacks. This endotype will include 3 of the newly described mutations—HAE-ANGPT, HAE-MYOF, and HAE-HSST (Table III)—and HAE-UNK, as well as systemic capillary leak syndrome (SCLS, also known as Clarkson disease).

Although there is little information on the underlying pathogenesis of HAE due to heparan sulfate 3-O-sulfotransferase 6 mutation (HAE-HSST), with only one family reported,²⁹ the disease-driving mutation affects heparin sulfate (HS)-glucosamine 3-O-sulfotransferase 6 (HS3ST6), a transmembrane enzyme involved in the synthesis of HS, part of the vascular endothelium infrastructure. HS contributes to the binding of high-molecular-weight kininogen (HK), which then becomes prone to cleavage by kallikrein.²⁹ Although some experts proposed

Categorization of HAE




FIG 3. Categorization of HAE.

TABLE III. Consented abbreviations/acronyms, in alphabetic order

Terminology in use	Acronym
Angioedema	AE
Bradykinin-mediated angioedema	AE-BK
Acquired angioedema due to <i>C1INH</i> deficiency	AAE-C1INH
Episodic angioedema with eosinophilia	EAE
Drug-induced angioedema	AE-DI
Mast cell-mediated angioedema	AE-MC
Angioedema due to unknown cause	AE-UNK
Angioedema due to vascular endothelium dysfunction	AE-VE
Angioedema due to anaphylaxis	AE-ANA
Angioedema due to normal <i>C1INH</i>	HAE-nC1INH
Chronic spontaneous urticaria	CSU
Hereditary angioedema	HAE
HAE due to angiopoietin 1 mutation	HAE-ANGPT
HAE due to <i>C1INH</i> deficiency	HAE-C1INH
HAE due to <i>C1INH</i> deficiency type 1	HAE-C1INH-Type1
HAE due to <i>C1INH</i> deficiency type 2	HAE-C1INH-Type2
HAE due to factor XII mutation	HAE-FXII
HAE due to heparan sulfate 3- <i>O</i> -sulfotransferase 6 mutation	HAE-HSST
HAE due to kininogen 1 mutation	HAE-KNG
HAE due to myoferlin mutation	HAE-MYOF
HAE due to plasminogen mutation	HAE-PLG
AE due to nonsteroidal anti-inflammatory drugs	AE-NSAID
AE due to angiotensin-converting enzyme inhibitors	AE-ACEI
AE due to tissue-type plasminogen activators	AE-tPA
AE due to di-peptidyl peptidase IV blockers	AE-DPPIV
AE due to angiotensin receptor–neprilysin inhibitors	AE-ARNI
Systemic capillary leak syndrome	SCLS
Urticaria	URT

Listed are drug families, abbreviations/acronyms, and generic names of drugs in common use implicated in acute and recurrent angioedema attacks.

considering HAE-HSST as BK mediated, the available evidence supports that it is due mainly to vascular endothelium dysfunction (Table I, statement 15).

Systemic capillary leak syndrome is a potentially fatal disorder characterized by unpredicted episodes of extensive AE, causing severe hypotension, hypoalbuminemia, renal failure, muscle necrosis due to compartment syndrome, and life-threatening multisystem failure.⁴⁷⁻⁴⁹ The pathogenesis of SCLS is held to be driven by endothelial cell dysfunction and subsequent plasma leakage, possibly involving vascular endothelial growth factor, angiopoietin 2, and certain interleukins.⁴⁹ Some experts argued that SCLS should not be classified as AE because it often manifests as generalized swelling. However, the majority of the panel voted to include it in the differential diagnosis of AE because it shares similar mechanisms (ie, vascular permeability) and several clinical features similar to HAE.

Drug-induced AE

A new category was created to include all drug-induced AE (AE-DI) because certain drugs can cause severe AE attacks, some of which may be fatal (Table I, statement 16).⁵⁰ The underlying pathomechanisms of most drugs listed in this category are unknown; therefore, establishing this distinct category is primarily practical. It aims to help clinicians differentiate AE due to adverse drug reactions (iatrogenic) from other types of AE and thereby avoid further exposure of the patients to the culprit drug or to drugs with similar class effects.

The AE-DI category includes ACE inhibitors (ACEI) and other inhibitors of the renin–angiotensin aldosterone system, including dipeptidyl peptidase IV (DPPIV) and angiotensin receptor–neprilysin inhibitors (ARNI), known triggers for acute drug-induced AE episodes in patients with hypertension, cardiac

disease, and/or diabetes (Table I, statement 17).⁵¹⁻⁶⁵ Some expert panel members argued that ACEI-induced AE should be designated AE-BK because ACE (kininase II) degrades BK, and BK levels are increased in patients with AE-ACEI. Similarly, DPPIV inhibitors degrade substance P when ACE is inhibited, increasing the risk of AE in individuals receiving ACEI therapy.⁵¹⁻⁵⁴ However, the majority of the panel voted for AE-DI because despite the empirical use of BK receptor antagonist icatibant, it failed to show efficacy in some of the randomized, placebo-controlled trials.^{54,55}

Nonsteroidal anti-inflammatory drugs (NSAIDs)⁶⁶ and inhibitors of tissue-type plasminogen activator (tPAs) are also known culprits for drug-induced orolingual AE.⁶⁷⁻⁷⁰ Recombinant tPA is used extensively in ischemic stroke and coronary thrombolysis. The mechanism of AE is still not fully understood, but new data suggest that tPA may trigger HK cleavage and induce BK formation.⁷¹ NSAIDs may trigger AE via eicosanoid pathways or via MC degranulation. Table IV lists pharmacologic agents implicated in acute AE.

AE of unknown cause

AE subtypes for which the pathomechanism or inciting trigger is unknown are classified as AE-UNK (Table I, statement 18). AE-UNK should be differentiated from HAE-UNK; the former is not defined by family history or other consensus clinical criteria, whereas the latter is.²⁰⁻²²

The inclusion of a category comprising unknown AE entities reflects an unmet diagnostic need that should inspire further research and yield new findings to enrich the current knowledge base. The DANCE experts recommend that patients with disease diagnosed in this category be followed up and provided the best of care, including off-label therapies judged indispensable by AE experts.

AE-UNK includes episodic AE with eosinophilia (EAE) (Table I, statement 19), a rare disorder characterized by unpredictable attacks of AE and high blood eosinophil levels.⁷² Clinical findings include recurrent or nonrecurrent AE, fever, URT, weight gain, blood eosinophilia, and elevated immunoglobulin levels.^{73,74} Some evidence points to a contribution of MC to its pathogenesis,⁷³ or to increased numbers of MCs and serum levels of tryptase. Eosinophil granule proteins were shown (in hamsters) to cause increased vasopermeability independently of MC and histamine, and other cell types, including neutrophils and lymphocytes, might be involved in pathogenesis.⁷³ Panel deliberations concluded that there is insufficient evidence to support any of these mechanisms, leading to the designation of EAE as AE-UNK.

DISCUSSION

Disease classification systems are essential to introduce order in conceptualizing human health disorders. Nosologic classifications disclose common disease pathophysiology and shared genetic associations, identify new drug targets, and classify disease for diagnostic, prognostic, and treatment decision purposes.⁷⁵ Disease classifications should bring on board emerging pathophysiologic, biochemical, and genomic data, making them widely available and easily shared by research networks.⁷⁵ Future machine learning may help to increase our understanding of HAE etiology, making it accessible for future applications of personalized medicine.⁷⁵

All AEs covered by the proposed classification are phenotypically expressed by recurrent, localized, and reversible episodes

of tissue swelling, regardless of the underlying mechanisms or triggers initiating the attacks. In all types and subtypes of AE, swelling is due to increased extravasation, in which the underlying pathophysiologic target is the vascular endothelium, whose continuity and integrity are breached by internal (ie, BK, histamine) and/or external stimuli (ie, drugs).⁸ A large body of evidence supports the notion that such a breach in functional continuity occurs within paracellular mechanisms of the cadherin junction.³ In all the different AEs, the protein-rich plasma entering the tissues is gradually evacuated after each attack by the local lymphatic system.^{3,4} Because AE types are different in their mechanisms and triggering signals, the DANCE classification takes these differences into account and aims to create a unified system covering all currently known AE entities. Such inclusive taxonomy will better assist clinicians in diagnosis and treatment decisions and guide basic research investigators in their efforts to elucidate the mechanisms of unknown diseases.

The DANCE expert panel recognize that AEs are not dichotomous and may have overlapping signatures, so we propose applying the term “endotype” to certain AE types. An endotype is a subtype defined by a distinct functional or pathobiologic mechanism to form a combination of clinical expression, biomarkers, and genetics.^{76,77} Such an approach has already proven its usefulness in chronic rhinitis with or without nasal polyposis³ and in asthma.^{78,79}

On the basis of this rationale, the DANCE experts propose to classify MC-mediated AE (AE-MC), BK-mediated AE (AE-BK), and vascular endothelium dysfunction AE (AE-VE) as endotypes, a designation that better reflects their underlying mechanisms, the mediators involved, and the results of genetic cosegregation studies, thereby justifying the existence of a distinct pathobiologic entity. We hope this approach helps focus future research in developing targeted treatments and applying personalized medicine to patients.

Less defined conditions, with mechanisms that remain to be elucidated, will be designated as subtypes. This applies in particular to those caused or aggravated by medications (henceforth AE-DI; ie, iatrogenic, drug induced), or AEs with still unknown etiologies (henceforth AE-UNK; ie, idiopathic, unknown). Despite gaps in our knowledge, organizing classes and subclasses of AEs by their underlying mechanisms, although sometimes presumptive, may lead to more accurate diagnosis and better selection of targeted pharmacotherapies.

The most challenging clinical dilemma among clinicians in emergency departments and daily practice is assessing whether a patient's swelling episode is due to MC mediators or BK. This common diagnostic challenge is particularly important when a decision concerning future treatment needs to be made. AE induced by MC mediators is by far the most frequently observed condition, as in CSU. In the absence of specific biomarkers, AE in CSU without wheals, sometimes referred to as MC-mediated AE, is difficult to diagnose and classify. One approach to the current differential diagnosis excludes other rare causes of recurrent AE and assesses response to antihistamines, corticosteroids, and omalizumab. Some authors argue that patients with AE in CSU without wheals should not be considered to have CSU due to clinical and laboratory differences compared to CSU patients with both recurrent wheals and AE.⁸⁰

Ongoing research efforts aim to identify biomarkers that distinguish AE attributed to the AE-MC endotype from those

TABLE IV. Drugs implicated in angioedema

Drug or drug family	Acronym	Examples	References
Angiotensin-converting enzyme inhibitors (-prils)	ACEI	Enalapril, ramipril, captopril, lisinopril, benazepril, cilazapril	50-60
Angiotensin receptor blockers (-sartans)*	ARB	Losartan, olmesartan, candesartan, telmisartan	51-54
Nonsteroidal anti-inflammatory drugs	NSAID	Aspirin, ibuprofen, naproxen	66
Recombinant human tissue plasminogen activator	rhtPA	Streptokinase, alteplase, tenecteplase	67-71
Dipeptidyl peptidase IV blockers	DPPIV	Sitagliptin, vildagliptin, saxagliptin	62,63
Angiotensin receptor–neprilysin inhibitors	ARNI	Sacubitril	64,65

*ACEI-induced angioedema is not a contraindication to angiotensin receptor blocker therapy.

mediated by other types of AE.^{81,82} Additionally, diagnostic algorithms may help establish a correct diagnosis by carefully evaluating the nature of clinical symptoms, family history, response to medications, and laboratory tests, thereby providing clues for clinical diagnosis and management of AE entities.⁸³

The global panel of experts agrees that episodic and recurrent AE caused by an increased vascular endothelial permeability deserves to be recognized as a separate nosologic classification. Unified classification, terminology, and nomenclature, along with a widely accepted list of abbreviations/acronyms, are necessary for better diagnosis, clinical decision making, and evidence-based therapeutic strategies. This approach can be challenging, thanks to the significant differences across the globe regarding access to advanced therapeutics, insurance payor systems, and other barriers to diagnosis and care that vary from one country to another. Nonetheless, it is imperative for the development of advanced treatment modalities through clinical trials and future personalized medicine that an agreed-on global terminology for AE disorders be established. Table III and Fig 2 summarize the updated changes in nomenclature and terminology of AE.

This proposed nomenclature system includes AE-UNK because many AE cases still lack substantive data regarding biomarkers, pathomechanisms, or genetics. The DANCE experts are optimistic that additional biomarkers and diagnostics evidence will be available in the future, contributing to a better understanding of AE-UNK. Therefore, we recommend regularly updating this consensus as new evidence on AE types and subtypes becomes available. It is essential to emphasize that the DANCE classification be used in complement with current clinical guidelines and expert consensus recommendations for diagnostic assessment and treatment of AE. They are not meant to replace current clinical guidelines and expert consensus algorithms, and they should not be misconstrued in a way that affects the reimbursement of medicines prescribed by a physician using sound clinical judgment.^{20-22,84}

In summary, DANCE has unified the definition, classification, nomenclature, and terminology of AE syndromes, and their use will harmonize and facilitate research, clinical studies, and the development of new medications, which will, we hope, result in improved patient care.

DISCLOSURE STATEMENT

The US HAE Association and HAE International provided unrestricted grants to support the DANCE project.

Disclosure of potential conflict of interest: A. Reshef has received speaker/advisor fees and/or research funding from SCL Behring, Ionis, Pharvaris, BioCryst, Teva, SIS Shulov, and

Takeda. T. Buttgereit has been speaker and/or advisor for AstraZeneca, BioCryst, CSL-Behring, GSK, Hexal, KalVista, Medac, Novartis, Pharming, Roche, Sanofi, and Takeda. A. S. Grumach has received speaker/advisor fees and/or research funding from CSL Behring, Pharvaris, KalVista, Biomarin, Pintpharma, Multicare, and Takeda. H. Farkas has received research grants from CSL Behring, Takeda, and Pharming; and has been an advisor for these companies as well as KalVista, ONO Pharmaceutical, Pharvaris, ASTRIA, Intellia, and BioCryst. H. Longhurst has received speaker/advisor fees and/or research funding from BioCryst, CSL Behring, Intellia, KalVista, Pharming, Pharvaris, and Takeda. J. Peter has received speaker/advisor fees and/or research funding from Astria, BioCryst, CSL Behring, Glenmark, Novartis, Pharming, Pharvaris, Sanofi/Regeneron, and Takeda. M. A. Riedl has received research support from BioCryst, Biomarin, CSL Behring, Intellia, Ionis, KalVista, Pharvaris, and Takeda; consulting fees from Astria, BioCryst, Biomarin, CSL Behring, Cycle, Intellia, KalVista, Ono Pharma, Pharming, Pharvaris, and Takeda; and honoraria for speaker presentations from CSL Behring, Pharming, and Takeda. M. Hide has received speaker/advisor fees and/or research funding from Amgen, Eisai, CSL-Behring, Kaken, KalVista, Kyorin Pharmaceutical, Kyowa-Kirin, Mitsubishi-Tanabe Pharma, Novartis, Pharvaris, Sanofi, Taiho, Takeda, Teikoku Seiyaku, and Torii Pharmaceutical. S. D. Betschel has received speaker/advisor fees and/or research funding from Astria, Canadian Blood Services, CSL Behring, Green Cross, Grifols, Ionis Pharmaceuticals, KalVista Novartis, Octapharma, Pharvaris, Sanofi, Takeda, and WSIB. T. Caballero has received speaker/advisor fees and/or research funding from Astria, BioCryst, CSL Behring, Ionis, KalVista, Novartis, Pharming NV, Pharvaris, and Takeda; and is researcher with IdiPAZ Program for promoting research activities. Y. Zhi has received speaker/advisor fees and/or research funding from Takeda. W. Aberer has received speaker/advisor fees and/or research funding from CSL Behring, BioCryst, and Takeda. S. Altrichter has received speaker/advisor fees and/or research funding from AstraZeneca, BioCryst, CSL Behring, LEO Pharma, Moxie GmbH, Novartis, Pharvaris, Sanofi, Takeda, and Thermo Fisher. E. Aygören-Pürsün has received speaker/advisor fees and/or research funding from Shire/Takeda, CSL Behring, Pharming, Biomarin, Centogene, KalVista, Pharvaris, BioCryst, and Intellia. M. L. Baeza has received speaker/advisor fees and/or research funding from CSL Behring, Takeda, Novartis, and BioCryst. N. A. Bara has received speaker/advisor fees and/or has performed clinical trials and/or received research funding from AbbVie, Almirall, Amgen, AstraZeneca, BioCryst, Biofrontera, Celldex, Centogene, CSL Behring, LaRoche Posay, Leo Pharma, Lilly, Novartis, Pfizer, Pharvaris, Sanofi/Regeneron, and Takeda. A. Bauer has received speaker/advisor fees and/or

research funding from Shire/Takeda, Pharming Group, and GlaxoSmithKline. J. A. Bernstein has received speaker/advisor fees and/or research funding from CSL Behring, Biomarin, Astria, Escent, Amgen, Celldex, Blueprint Medicines, Jasper, Teva, Takeda/Shire, BioCryst, Pharming, Ionis, KalVista, Pharvaris, Novartis, Sanofi/Regeneron, Genentech, Celldex, Allakos, and AstraZeneca. I. Boccon-Gibod has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Pharming, Novartis, and BioCryst. H. Bonnekoh has received speaker/advisor fees and/or research funding from AbbVie, Intercept Pharma, Novartis, Sanofi Aventis, and Valenza Bio. L. Bouillet has received speaker/advisor fees, research and educational funding, and/or travel grants from BioCryst, CSL Behring, Takeda, Novartis, GSK, and Blueprint Medicines. A. Bygum has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Viropharma, and Novartis. O. Calderon has received speaker/advisor fees and/or research funding from Pint Pharma. R. de Albuquerque Campos has received speaker/advisor fees and/or research funding from Novartis and Takeda. M. Cancian has received speaker/advisor fees and/or research funding from Shire/Takeda, CSL Behring, Pharming, SOBI, and BioCryst. H. J. Chong-Neto has received speaker/advisor fees and/or research funding from Takeda, AstraZeneca, Viatrix, Sanofi, Abbott, and Thermo Fisher. S. Cimbollek has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, and Novartis. D. M. Cohn has received speaker/advisor fees and/or research funding from CSL Behring, Takeda, Pharvaris, BioCryst, Ionis Pharmaceuticals, KalVista, and Pharming. T. Craig has been speaker and researcher for Biomarin, CSL-Behring, Intellia, and Takeda; researcher for Astria, Ionis, KalVista, and Pharvaris; speaker for Grifols; consultant for Astria, BioCryst, Biomarin, CSL Behring, Intellia, and Spark; director for ACARE International Hereditary Angioedema Center; and member of medical advisory board for HAE-A. I. Danilycheva has received speaker/advisor fees and/or research funding from Novartis, Generium Pharmaceutical, and Berlin-Chemie AG/Menarini Group. A. Du-Thanh has received speaker/advisor fees and/or research funding from Shire/Takeda and Novartis. L. F. Ensina has received speaker/advisor fees and/or research funding from AbbVie, Novartis, and Sanofi. D. Fomina has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Novartis, Sanofi, Roche, and AstraZeneca. L. Fonacier has received advisor fees and/or research funding from Regeneron, Pfizer, AstraZeneca, AbbVie, Lilly, and Leo. A. Fukunaga has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Torii Pharmaceutical, Novartis, Tanabe Pharmaceutical, Taiho Pharmaceutical, Kaken Pharmaceutical, and Kyorin Pharmaceutical. A. Gelincik has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Vem Ilac, and Polifarma. P. Giavina-Bianchi has received speaker/advisor fees and/or research funding from Takeda and CSL Behring. M. Goncalo has received speaker/advisor fees and/or research funding from Takeda, Leo Pharma, Lilly, Pfizer, AbbVie, Sanofi, and Novartis. M. Guilarte has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Novartis, Pharvaris, KalVista, BioCryst, and BioMarin. T. Kinacyan has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, BioCryst, KalVista, Pharvaris, Sanofi, Novartis, and Kiniksa. P. Kolkhir has received speaker/advisor fees and/or research funding from Novartis and Roche. K.

Kulthanan has received speaker/advisor fees and/or research funding from A. Menarini and Novartis. M. Kurowski has received speaker/advisor fees and/or research funding from CSL Behring, Pharming, Takeda, and KalVista. E. Latysheva has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Generium Pharmaceutical, and Octapharma. D. Launay has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, BioCryst, Octapharma, Boehringer Ingelheim, Mallinckrodt Pharmaceuticals, and Blueprint Medicines. R. Lleonhart has received speaker/consultancy fees from BioCryst, CSL-Behring, KalVista Novartis, and Takeda; has received funding to attend conferences/educational events from CSL Behring, Novartis, Pharming, and Takeda; and has participated as investigator in clinical trial/registry for BioCryst, Takeda, KalVista, Pharvaris, and IONIS. W. Lumry has received speaker/advisor fees and/or research funding from AstraZeneca, Astria, BioCryst, Biomarin, CSL Behring, GSK, Ionis, Intellia, KalVista, Pharming, Phavaris, Sanofi/Regeneron, and Shire/Takeda. R. M. Ali has been speaker for Takeda. S. Nieto-Martinez has been speaker for Takeda. G. Porebski has received speaker/advisor fees and/or research funding from CSL Behring, Pharming, and Takeda. A. Prieto Garcia has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, Pharvaris, and Bio-Cryst. F. S. Serpa has received speaker/advisor fees from Takeda and CSL Behring. F. Siebenhaar has received speaker/advisor fees and/or research funding from Allakos, Blueprint Medicines, Celldex, Cogent, Escent, Granular, GSK, Invea, Noucor, Novartis, Moxie, Sanofi/Regeneron, and ThirdHarmonicBio. M. Staevska has received speaker/advisor fees and/or research funding from Shire/Takeda, Pharming Group, CSL Behring, SOBI, Ionis, Pharvaris, KalVista, AstraZeneca, Sanofi, Novartis, AbbVie, GSK, Menarini, Organon, and Chiesi. P. Staubach has received speaker/advisor fees and/or research funding from Takeda, Viropharm, CSL Behring, Sanofi, Novartis, Octapharma, and Pharming. M. Stobiecki has received speaker/advisor fees and/or research funding from Pharming, CSL Behring, KalVista, BioCryst, Takeda, Astria, and Pharming. S. F. Thomsen has received speaker/advisor fees and/or research funding from AbbVie, Janssen, LEO Pharma, Novartis, Sanofi, UCB, Eli Lilly, and Pfizer. A. Valerieva has received speaker/advisor fees and/or research funding from Shire/Takeda, Pharming Group, CSL Behring, SOBI, Pharvaris, KalVista, Ionis Pharmaceuticals, Astria Therapeutics, BioCryst, and Novartis. N. Van Dinh has received speaker/advisor fees and/or research funding from Takeda, CSL Behring, and Novartis. A. Zalewska-Janowska has received advisor fees from NAOS. A. Zanichelli has received speaker/advisor fees and/or research funding from Astria, CSL Behring, BioCryst, KalVista, Takeda, and Pharming. M. Magerl has received honoraria and/or consultation fees from Takeda, CSL Behring, KalVista, Pharvaris, and Astria. M. Maurer has received speaker/advisor fees and/or research funding from Allakos, Alvotech, Amgen, Aquestive, Aralez, AstraZeneca, Astria, Bayer, BioCryst, Blueprint Medicines, Celldex, Celltrion, Centogene, CSL Behring, Evoimmune, GSK, Ipsen, KalVista, Kyowa Kirin, Leo Pharma, Lilly, Menarini, Mitsubishi Tanabe Pharma, Moxie, Noucor, Novartis, Orion Biotechnology, Pharvaris, Resonance Medicine, Sanofi/Regeneron, Septerna, Takeda, Teva, Third HarmonicBio, Valenza Bio, Yuhan Corporation, and Zurabio. The rest of the authors declare that they have no relevant conflicts of interest.

The DANCE project benefited from the network of ACAREs of GA²LEN, the Global Allergy and Asthma Excellence Network. We thank Carolina Vera at the Berlin ACARE for her help with reviewing the literature on AE. We are thankful to ACARE, ACAAI, APAAACI, and AAAAI for their endorsement of the initiative, the nomination of the panel members, and their commitment for the dissemination of the DANCE classification and scientific vocabulary.

Key messages

- Up-to-date comprehensive classification and terminology covering a wide range of AE syndromes was consented by an international team.
- This terminology will facilitate diagnosis, research, new medication development, and patient care.

REFERENCES

- Cicardi M, Aberer W, Banerji A, Bas M, Bernstein JA, Bork K, et al. Classification, diagnosis, and approach to treatment for angioedema: consensus report from the Hereditary Angioedema International Working Group. *Allergy* 2014;69:602-16.
- Wu MA, Perego F, Zanichelli A, Cicardi M. Angioedema phenotypes: disease expression and classification. *Clin Rev Allergy Immunol* 2016;51:162-9.
- Claesson-Welsh L, Dejana E, McDonald DM. Permeability of the endothelial barrier: identifying and reconciling controversies. *Trends Mol Med* 2021;27:314-31.
- Busse PJ, Christiansen SC. Hereditary angioedema. *N Engl J Med* 2020;382:1136-48.
- Mormile I, Cocchiari A, Bova M, Loffredo S, de Paulis A, Spadaro G, et al. Gastrointestinal manifestations of angioedema: a potential area of misdiagnosis. *Eur J Gastroenterol Hepatol* 2021;33(6):787-93.
- Zuberbier T, Abdul Latiff AH, Abuzakouk M, Aquilina S, Asero R, Baker D, et al. The international EAACI/GA²LEN/EuroGuiDerm/APAAACI guideline for the definition, classification, diagnosis, and management of urticaria. *Allergy* 2022;77:734-66.
- Maurer M, Magerl M. Differences and similarities in the mechanisms and clinical expression of bradykinin-mediated vs mast cell-mediated angioedema. *Clin Rev Allergy Immunol* 2021;61:40-9.
- Kaplan AP, Joseph K. Pathogenesis of hereditary angioedema: the role of the bradykinin-forming cascade. *Immunol Allergy Clin N Am* 2017;37:513-25.
- Donaldson VH, Evans RR. A biochemical abnormality in hereditary angioneurotic edema: absence of serum inhibitor of C' 1-esterase. *Am J Med* 1963;35:37-44.
- Landerman NS. Hereditary angioneurotic edema. *J Allergy* 1962;33:116-29.
- Germenis AE, Margaglione M, Pesquero JB, Farkas H, Cichon S, Csuka D, et al. International consensus on the use of genetics in the management of hereditary angioedema. Hereditary Angioedema International Working Group. *J Allergy Clin Immunol Pract* 2020;8:901-11.
- Ponard D, Gaboriaud C, Charignon D, Ghannam A, Wagenaar-Bos IGA, Roem D, et al. *SERPINI1* mutation update: mutation spectrum and C1 inhibitor phenotypes. *Hum Mutat* 2020;41:38-57.
- Germenis AE, Rijavec M, Veronez CL. Leveraging genetics for hereditary angioedema: a road map to precision medicine. *Clin Rev Allergy Immunol* 2021;60:416-28.
- Bork K, Hardt J, Witzke G. Fatal laryngeal attacks and mortality in hereditary angioedema due to C1-INH deficiency. *J Allergy Clin Immunol* 2012;130:692-7.
- Minafra FG, Gonçalves TR, Alves TM, Pinto JA. The mortality from hereditary angioedema worldwide: a review of the real-world data literature. *Clin Rev Allergy Immunol* 2022;62:232-9.
- Longhurst H, Bygum A. The humanistic, societal, and pharmaco-economic burden of angioedema. *Clin Rev Allergy Immunol* 2016;51:230-9.
- Caballero T, Prior N. Burden of illness and quality-of-life measures in angioedema conditions. *Immunol Allergy Clin North Am* 2017;37:597-616.
- Lumry WR, Settupane RA. Hereditary angioedema: epidemiology and burden of disease. *Allergy Asthma Proc* 2020;41(Suppl 1):S08-13.
- Banerji A, Davis KH, Brown TM, Hollis K, Hunter SM, Long J, et al. Patient-reported burden of hereditary angioedema: findings from a patient survey in the United States. *Ann Allergy Asthma Immunol* 2020;124:600-7.
- Maurer M, Magerl M, Betschel S, Aberer W, Ansotegui IJ, Aygoren-Pursun E, et al. The international WAO/EAACI guideline for the management of hereditary angioedema—the 2021 revision and update. *Allergy* 2022;77:1961-90.
- Betschel S, Badiou J, Binkley K, Borici-Mazi R, Hebert J, Kanani A, et al. The international/Canadian hereditary angioedema guideline. *Allergy Asthma Clin Immunol* 2019;15:1-29.
- Busse PJ, Christiansen SC, Riedl MA, Banerji A, Bernstein JA, Castaldo AJ, et al. US HAEA medical advisory board 2020 guidelines for the management of hereditary angioedema. *J Allergy Clin Immunol Pract* 2021;9:132-50.e3.
- Bork K, Barnstedt SE, Koch P, Traupe H. Hereditary angioedema with normal C1-inhibitor activity in women. *Lancet* 2000;356(9225):213-7.
- Binkley KE, Davis A III. Clinical, biochemical, and genetic characterization of a novel estrogen-dependent inherited form of angioedema. *J Allergy Clin Immunol* 2000;106:546-50.
- Bork K, Wulff K, Steinmüller-Magin L, Brænne I, Staubach-Renz P, Witzke G, et al. Hereditary angioedema with a mutation in the plasminogen gene. *Allergy* 2018;73:442-50.
- Bafunno V, Firinu D, D'Apolito M, Cordisco G, Loffredo S, Leccese A, et al. Mutation of the angiotensin-converting enzyme 1 gene (*ANGPT1*) associates with a new type of hereditary angioedema. *J Allergy Clin Immunol* 2018;141:1009-17.
- Bork K, Wulff K, Rossmann H, Steinmüller-Magin L, Brænne I, Witzke G, et al. Hereditary angioedema co-segregating with a novel kininogen 1 gene mutation changing the N-terminal cleavage site of bradykinin. *Allergy* 2019;74:2479-81.
- Ariano A, D'Apolito M, Bova M, Bellanti F, Loffredo S, D'Andrea G, et al. A myoferlin gain-of-function variant associates with a new type of hereditary angioedema. *Allergy* 2020;75:2989-92.
- Bork K, Wulff K, Möhl BS, Steinmüller-Margin L, Witzke G, Hardt J, et al. Novel hereditary angioedema linked with a heparan sulfate 3-O-sulfotransferase 6 gene mutation. *J Allergy Clin Immunol* 2021;148:1041-8.
- Veronez CL, Csuka D, Sheikh FR, Zuraw BL, Farkas H, Bork K. The expanding spectrum of mutations in hereditary angioedema. *J Allergy Clin Immunol Pract* 2021;9:2229-34.
- Reshef A, Kidon M, Leibovich I. The story of angioedema: from quincke to bradykinin. *Clin Rev Allergy Immunol* 2016;51:121-39.
- Reshef A. Hereditary angioedema: a call for collective terminology. *Allergy Asthma Proc* 2016;37:e14.
- Banerji A. Hereditary angioedema: a call for collective terminology (response). *Allergy Asthma Proc* 2016;37:e14.
- Aberer W. Hereditary angioedema: an orphan but an original disease? *J Allergy Clin Immunol* 2021;148:994-5.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analysis: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006-12.
- Dalkey N. An experimental study of group opinion: the Delphi method. *Futures* 1969;1:408-26.
- Akins RB, Tolson H, Cole BR. Stability of response characteristics of a Delphi panel: application of bootstrap data expansion. *BMC Med Res Methodol* 2005;5:37.
- Diamond IR, Grant RC, Feldman BM, Pencharz PB, Ling SC, Moore AM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67:401-9.
- Maurer M, Aygoren-Pursun E, Banerji A, Bernstein JA, Boysen HB, Busse PJ, et al. Consensus on treatment goals in hereditary angioedema: a global Delphi initiative. *J Allergy Clin Immunol* 2021;148:1526-32.
- Weerasubpong P, Jiamton S, Phumariyapong P, Ungprasert P, Kulthanan K. Prevalence of concomitant angioedema in chronic spontaneous urticaria: a systematic review and meta-analysis. *Asian Pac J Allergy Immunol* 2023;41:12-9.
- Leibovich-Nassi I, Reshef A. The enigma of prodromes in hereditary angioedema (HAE). *Clin Rev Allergy Immunol* 2021;61:15-28.
- Magerl M, Doumoulakis G, Kalkounou I, Weller K, Church MK, Kreuz W, et al. Characterization of prodromal symptoms in a large population of patients with hereditary angio-oedema. *Clin Exp Dermatol* 2014;39:298-303.
- Schmaier AH, Emsley J, Feener EP, Gailani D, Govers-Riemsag JWP, Kaplan AP, et al. Nomenclature of factor XI and the contact system. *J Thromb Haemost* 2019;17:2216-9.
- Galvan-Blasco P, Gil-Serrano J, Sala-Cunill A. New biomarkers in anaphylaxis (beyond tryptase). *Curr Treat Options Allergy* 2022;9:303-22.
- Muraro A, Worm M, Alviani C, Cardona V, DunnGalvin A, Garvey LH, et al. EAACI guidelines: anaphylaxis (2021 update). *Allergy* 2022;77:357-77.
- Hintze S, Möhl BS, Beyerl J, Wulff K, Wieser A, Bork K, Meinke P. Mutant plasminogen in hereditary angioedema is bypassing FXII/kallikrein to generate bradykinin. *Front Physiol* 2023;13:1090732.
- Clarkson B, Thompson D, Horwith M, Luckey EH. Cyclical edema and shock due to increased capillary permeability. *Am J Med* 1960;29:193-216.
- Xie Z, Chan E, Yin Y, Ghosh CC, Wisch L, Nelson C, et al. Inflammatory markers of the systemic capillary leak syndrome (Clarkson disease). *J Clin Cell Immunol* 2014;5:1000213.

49. Eo TS, Chun KJ, Hong SJ, Kim JY, Lee IR, Lee KH, et al. Clinical presentation, management, and prognostic factors of idiopathic systemic capillary leak syndrome: a systematic review. *J Allergy Clin Immunol Pract* 2018;6:609-18.
50. Lochbaum R, Hoffmann TK, Greve J, Hahn J. Concomitant medication in patients with bradykinin-mediated angioedema—there's more than ACE inhibitors. *J Dtsch Dermatol Ges* 2023;21:1283-9.
51. Stone C Jr, Brown NJ. Angiotensin-converting enzyme inhibitor and other drug-associated angioedema. *Immunol Allergy Clin North Am* 2017;37:483-95.
52. Brown T, Gonzalez J, Monteleone CJ. Angiotensin-converting enzyme inhibitor-induced angioedema: a review of the literature. *Clin Hypertens (Greenwich)* 2017;19:1377-82.
53. Kostis WJ, Shetty M, Chowdhury YS, Kostis JB. ACE inhibitor-induced angioedema: a review. *Curr Hypertens Rep* 2018;20:55.
54. Rasmussen ER, Hallberg P, Baranova EV, Eriksson N, Karawajczyk M, Johansson C, et al. Genome-wide association study of angioedema induced by angiotensin-converting enzyme inhibitor and angiotensin receptor blocker treatment. *Pharmacogenomics J* 2020;20:770-83.
55. Montinaro V, Cicardi M. ACE inhibitor-mediated angioedema. *Int Immunopharmacol* 2020;78:106081.
56. Sinert R, Levy P, Bernstein JA, Body R, Sivilotti MLA, Moellman J, et al. Randomized trial of icaltiban for angiotensin-converting enzyme inhibitor-induced upper airway angioedema. *J Allergy Clin Immunol Pract* 2017;5:1402-9.
57. Straka BT, Ramirez CE, Byrd JB, Stone E, Woodard-Grice A, Nian H, et al. Effect of bradykinin receptor antagonism on ACE inhibitor-associated angioedema. *J Allergy Clin Immunol* 2017;140:242-8.e2.
58. Bas M, Greve J, Strassen U, Khosravani F, Hoffmann TK, Kojda G. Angioedema induced by cardiovascular drugs: new players join old friends. *Allergy* 2015;70:1196-2000.
59. Banerji A, Blumenthal KG, Lai KH, Zhou L. Epidemiology of ACE inhibitor angioedema utilizing a large electronic health record. *J Allergy Clin Immunol Pract* 2017;5:744-9.
60. Weisman DS, Arnouk N, Asghar MB, Qureshi MR, Kumar A, Desale S, et al. ACE inhibitor angioedema: characterization and treatment versus non-ACE angioedema in acute hospitalized patients. *J Community Hosp Intern Med Perspect* 2020;10:16-8.
61. Leru PM, Anton VF, Bocsan C, Muntean A, Boda D. Acquired angioedema induced by angiotensin-converting enzyme inhibitors—experience of a hospital-based allergy center. *Exp Ther Med* 2020;20:68-72.
62. Byrd JB, Shreevatsa A, Putlur P, Foretia D, McAlexander L, Sinha T, et al. Dipeptidyl peptidase IV deficiency increases susceptibility to angiotensin-converting enzyme inhibitor-induced peritracheal edema. *J Allergy Clin Immunol* 2007;120:403-8.
63. Cassano N, Nettis E, Di Leo E, Ambrogio F, Vena GA, Foti C. Angioedema associated with dipeptidyl peptidase-IV inhibitors. *Clin Mol Allergy* 2021;19:24.
64. Shi V, Senni M, Streefkerk H, Modgill V, Zhou W, Kaplan A. Angioedema in heart failure patients treated with sacubitril/valsartan (LCZ696) or enalapril in the PARADIGM-HF study. *Int J Cardiol* 2018;264:118-23.
65. Dani SS, Ganatra S, Vaduganathan M. Angioedema with sacubitril/valsartan: trial-level meta-analysis of over 14,000 patients and real-world evidence to date. *Int J Cardiol* 2021;323:188-91.
66. Kowalski ML, Woessner K, Sanak M. Approaches to the diagnosis and management of patients with a history of nonsteroidal anti-inflammatory drug-related urticaria and angioedema. *J Allergy Clin Immunol* 2015;136:245-51.
67. Myslimi F, Caparros F, Dequatre-Ponchelle N, Moulin S, Gautier S, Girardie P, et al. Orolingual angioedema during or after thrombolysis for cerebral ischemia. *Stroke* 2016;47:1825-30.
68. Szczepanski M, Bozyk P. Institutional incidence of severe tPA-induced angioedema in ischemic cerebral vascular accidents. *Crit Care Res Pract* 2018;2018:9360918.
69. Fröhlich K, Macha K, Gerner ST, Bobinger T, Schmidt M, Dörfler A, et al. Angioedema in stroke patients with thrombolysis. *Stroke* 2019;50:1682-7.
70. Vigneron C, Lécluse A, Ronzière T, Bouillet L, Boccon-Gibod I, Gayet S, et al. Angioedema associated with thrombolysis for ischemic stroke: analysis of a case-control study. *J Intern Med* 2019;286:702-10.
71. Charest-Morin X, Hebert J, Rivard G, Bonnefoy A, Wagner E, Marceau F. Comparing pathways of bradykinin formation in whole blood from healthy volunteers and patients with hereditary angioedema due to C1 inhibitor deficiency. *Front Immunol* 2018;9:2183.
72. Gleich GJ, Schroeter AL, Marcoux P, Sachs MI, O'Connell EJ, Kohler PF. Episodic angioedema associated with eosinophilia. *N Engl J Med* 1984;310:1621-6.
73. Khoury P, Herold J, Alpaugh A, Dinerman E, Holland-Thomas N, Stoddard J, et al. Episodic angioedema with eosinophilia (Gleich syndrome) is a multilineage cell cycling disorder. *Haematologica* 2015;100:300-7.
74. Haber R, Chebli JA, Gemayel EL, Salloum A. Gleich syndrome: a systematic review. *Int J Dermatol* 2020;59:1458-65.
75. Dozmorov MG. Disease classification: from phenotypic similarity to integrative genomics and beyond. *Brief Bioinform* 2019;20:1769-80.
76. Khurana Hershey GK, Sherenian MG, Mersha TB. Precision medicine. In: O'Hehir RE, Holgate ST, Khurana Hershey GK, Sheikh A, editors. *Allergy essentials*. 2nd ed. Amsterdam: Elsevier; 2021. pp. 25-39.
77. Agache I, Akdis CA. Precision medicine and phenotypes, endotypes, genotypes, regiotypes, and theratypes of allergic diseases. *J Clin Invest* 2019;129:1493-503.
78. Staudacher AG, Peters AT, Kato A, Stevens WW. Use of endotypes, phenotypes, and inflammatory markers to guide treatment decisions in chronic rhinosinusitis. *Ann Allergy Asthma Immunol* 2020;124:318-25.
79. Kaur R, Chupp G. Phenotypes and endotypes of adult asthma: moving toward precision medicine. *J Allergy Clin Immunol* 2019;144:1-12.
80. Sabaté-Brescó M, Rodríguez-Garijo N, Azofra J, Baeza ML, Donado CD, Gaig P, et al. A comparative study of sex distribution, autoimmunity, blood, and inflammatory parameters in chronic spontaneous urticaria with angioedema and chronic histaminergic angioedema. *Allergy Clin Immunol Pract* 2021;9:2284-92.
81. Lara-Marquez ML, Christiansen SC, Riedl MA, Herschbach J, Zuraw BL. Threshold-stimulated kallikrein activity distinguishes bradykinin- from histamine-mediated angioedema. *Clin Exp Allergy* 2018;48:1429-38.
82. Bindke G, Gehring M, Wiczorek D, Kapp A, Buhl T, Wedi B. Identification of novel biomarkers to distinguish bradykinin-mediated angioedema from mast cell-/histamine-mediated angioedema. *Allergy* 2022;77:946-55.
83. Farkas H, Balla Z, Riedl MA. Differentiating histaminergic and nonhistaminergic angioedema with or without urticaria. *J Allergy Clin Immunol* 2022;150:1405-9.
84. Zuraw BL, Bork K, Binkley KE, Banerji A, Christiansen SC, Castaldo A, et al. Hereditary angioedema with normal C1 inhibitor function: consensus of an international expert panel. *Allergy Asthma Proc* 2012;33(suppl 1):S145-56.

TABLE E1. Endorsing professional societies

No.	Acronym	Society
1	AAAAI	American Academy of Allergy, Asthma & Immunology
2	AAAeIC	Argentine Association of Allergy and Clinical Immunology
3	AAIBA	Association of Allergy and Immunology of Buenos Aires
4	ACAAI	American College of Allergy Asthma & Immunology
5	AID	Turkish National Society of Allergy and Clinical Immunology
6	ALaeh	Latin American Association of HAE
7	ALLSA	Allergy Society of South Africa
8	APAAACI	Asia Pacific Association of Allergy, Asthma and Clinical Immunology
9	ASBAI	Brazilian Association of Allergy and Immunology
10	ASCIA	Australasian Society of Clinical Immunology and Allergy
11	BACTA	Bulgarian Alliance of Clinical and Translational Allergy
12	BSA	Bulgarian Society of Allergology
13	CMICA	Mexican College of Clinical Immunology and Allergy
14	DDO	Danish Dermatology Society
15	DGAKI	German Society for Allergology and Clinical Immunology
16	DGIM	German Society of Internal Medicine
17	DST	Dermatology Society of Thailand
18	GA ² LEN	Global Allergy and Asthma European Network
19	GAACI	Georgian Association of Allergology and Clinical Immunology
20	GEBRAEH	Brazilian Study Group on Hereditary Angioedema
21	GSAR	German Society for Angioedema Research
22	HAESI	Hereditary Angioedema Society of India
23	HSACI	Hungarian Society for Allergy and Clinical Immunology
24	IAACI	Israeli Association of Allergy and Clinical Immunology
25	IADVL	Indian Association of Dermatologists, Venereologists & Leprologists
26	ITACA	Italian Network on Hereditary and Acquired Angioedema
27	JDA	Japanese Dermatological Association
28	KAAACI	Korean Academy of Asthma, Allergy, and Clinical Immunology
29	KSACI	Kuwait Society of Allergy and Clinical Immunology
30	MAKIT	Hungarian Society of Allergology and Clinical Immunology
31	NDO	Nordic Dermatology Association
32	NIV	Dutch Society for Internal Medicine
33	ÖGDV	Austrian Society of Dermatology and Venereology
34	OSACI	Omani Society of Allergy & Clinical Immunology
35	PDS	Polish Dermatological Society
36	PSA/PTA	Polish Society of Allergology
37	QAIS	Qatar Allergy and Immunology Society
38	RAACI	Russian Association of Allergology and Clinical Immunology
39	RSACI	Romanian Society of Allergology and Clinical Immunology
40	RUBRA	Rede Urticária Brazil
41	SBI	Brazilian Society of Immunology
42	SBP	Brazilian Society of Pediatrics
43	SEAIC	Spanish Society of Allergology and Clinical Immunology
44	SFA	French Society of Allergology
45	SFD	French Society of Dermatology
46	SIAAIC	Italian Society of Allergy, Asthma and Clinical Immunology
47	SIMI	Italian Society of Internal Medicine
48	SLaai	Latin American Society of Allergy and Immunology
49	SORESS	Society of Rhinology and Endoscopic Skull Base Surgery
50	SPAAI	Peruvian Society Allergy, Asthma and Immunology
51	SPDV	Portuguese Society of Dermatology and Venerology
52	TNSACI	Turkish National Society of Allergy and Clinical Immunology
53	UKPIN	United Kingdom Primary Immunodeficiency Network